

SUPER-SEQUENCES IN THE ARC COMPONENT OF A COMPACT CONNECTED GROUP

DIKRAN DIKRANJAN AND DMITRI SHAKHMATOV

Dedicated to Karl H. Hofmann on the occasion of his 76th anniversary

ABSTRACT. Let G be an abelian topological group. The symbol $\widehat{G}$ denotes the group of all continuous characters $\chi : G \rightarrow \mathbb{T}$ endowed with the compact open topology. A subset E of G is said to be *qc-dense* in G provided that $\chi(E) \subseteq \varphi([-1/4, 1/4])$ holds only for the trivial character $\chi \in \widehat{G}$, where $\varphi : \mathbb{R} \rightarrow \mathbb{T} = \mathbb{R}/\mathbb{Z}$ is the canonical homomorphism. A *super-sequence* is a non-empty compact Hausdorff space S with at most one non-isolated point (to which S converges). We prove that an infinite compact abelian group G is connected if and only if its arc component G_a contains a super-sequence converging to 0 that is qc-dense in G . This gives as a corollary a recent theorem of Außenhofer: For a connected locally compact abelian group G , the restriction homomorphism $r : \widehat{G} \rightarrow \widehat{G}_a$ defined by $r(\chi) = \chi|_{G_a}$ for $\chi \in \widehat{G}$, is a topological isomorphism. We also show that an infinite compact group G is connected if and only if its arc component G_a contains a super-sequence S converging to the identity e that generates a dense subgroup of G (equivalently, $S \setminus \{e\}$ is an infinite suitable set for G in the sense of Hofmann and Morris).

1. INTRODUCTION

All topological groups are assumed to be Hausdorff. Let G be a topological group. We denote by $\widehat{G}$ the group of all continuous characters $\chi : G \rightarrow \mathbb{T}$ endowed with the compact open topology. A subgroup D of G *determines* G if the restriction homomorphism $r : \widehat{G} \rightarrow \widehat{D}$ defined by $r(\chi) = \chi|_D$ for $\chi \in \widehat{G}$, is a topological isomorphism [5]. If G is locally compact and abelian, then every subgroup D that determines G must be dense in G . (When D is dense in G , the map $r : \widehat{G} \rightarrow \widehat{D}$ is a continuous isomorphism.) The following two theorems are the cornerstone results in the topic of determining subgroups:

Theorem 1.1. [1, 4] *A metrizable abelian group G is determined by each dense subgroup of G .*

Theorem 1.2. [12] *Every non-metrizable compact group G contains a dense subgroup that does not determine G .¹*

According to a well-known classical result of Eilenberg and Pontryagin, in a connected locally compact abelian group G the arc component G_a is dense. Since a subgroup of a

1991 *Mathematics Subject Classification.* Primary: 22C05; Secondary: 22B05, 54D05.

Key words and phrases. compact group, abelian group, connected space, pathwise connected space, arc component, super-sequence, suitable set, dual group.

The first author was partially supported by MEC. MTM2006-02036 and FEDER FUNDS.

The second author was partially supported by the Grant-in-Aid for Scientific Research no. 19540092 by the Japan Society for the Promotion of Science (JSPS).

¹A proof of this theorem under the assumption of CH can be found in [5].

locally compact abelian group determining it must be dense, the following theorem, recently proved by Außenhofer, is a strengthening of this classical result:

Theorem 1.3. [3] *The arc component G_a of a connected locally compact abelian group G determines G .*

While this theorem is a corollary of Theorem 1.1 for a metrizable group G , in the non-metrizable case the mere density of G_a in G ensured by the classical result of Eilenberg and Pontryagin mentioned above, need not guarantee that G_a determines G (as witnessed by Theorem 1.2).

Theorem 1.3 is used in [2] to prove that the uncountable powers of $\mathbb{Z}$ are not strongly reflexive, thereby resolving a problem raised by Banaszczyk on whether uncountable powers of the reals $\mathbb{R}$ are strongly reflexive.

Let $\varphi : \mathbb{R} \rightarrow \mathbb{T} = \mathbb{R}/\mathbb{Z}$ be the canonical homomorphism and $\mathbb{T}_+ = \varphi([-1/4, 1/4])$. We will say that a subset E of a topological group G is *qc-dense in G* (an abbreviation for *quasi-convexly dense*) provided that $\chi(E) \subseteq \mathbb{T}_+$ only for the trivial continuous homomorphism $\chi : G \rightarrow \mathbb{T}$.

This notion was introduced in [6] in the abelian context, and its significance for applications has been recently demonstrated in [10]. In particular, qc-density was used in [10] to establish essential properties of determining subgroups of compact abelian groups, thereby allowing to get a short elementary proof of Theorem 1.2.

The host of applications of qc-dense sets is made possible by the ultimate connection between the notions of determining subgroup and qc-density described in the next fact (proved in [10, Fact 1.4]). It is a particular case of a more general fact stated without proof (and in equivalent terms) in [5, Remark 1.2(a)] and [12, Corollary 2.2].

Fact 1.4. *A subgroup D of a compact abelian group G determines it if and only if there exists a compact subset of D that is qc-dense in G .*

It has been recently shown in [10] that qc-dense compact subsets (and thus determining subgroups) of a compact abelian group must be rather big.

Theorem 1.5. [10, Corollary 2.2] *If a closed subset X of an infinite compact abelian group G is qc-dense in G , then $w(X) = w(G)$. (Here $w(X)$ denotes the weight of a space X .)*

A *super-sequence* is a non-empty compact Hausdorff space X with at most one non-isolated point x^* [9]. We will call x^* the *limit* of X and say that X *converges to x^** . Observe that a countably infinite super-sequence is a convergent sequence (together with its limit).

Außenhofer [1] essentially proved that every infinite compact metric abelian group has a qc-dense sequence converging to 0.² This result has been recently extended to all compact groups by replacing convergent sequences with super-sequences:

Theorem 1.6. [10] *Every infinite compact abelian group contains a qc-dense super-sequence converging to 0.*

A subspace X of a topological group G *topologically generates G* if the subgroup of G generated by X is dense in G . The proof of the following fact is straightforward.

²This is an immediate consequence of [1, Theorem 4.3 or Corollary 4.4]. In fact, a more general statement immediately follows from these results: Every dense subgroup D of a compact metric abelian group G contains a sequence converging to 0 that is qc-dense in G .

Fact 1.7. [10, Fact 1.3(ii)] *Every qc-dense subset of a compact abelian group topologically generates it.*

Let G be a topological group with the identity e . If a discrete subset S of G topologically generates G and $S \cup \{e\}$ is closed in G , then S is called a *suitable set for G* [13].

Remark 1.8. Clearly, if S is a super-sequence in G that converges to e and topologically generates G , then $S \setminus \{e\}$ is a suitable set for G . Conversely, if G is compact and S is a suitable set for G , then $S \cup \{e\}$ must be a super-sequence.

It follows from this remark that a subgroup D of a compact group G contains a super-sequence converging to the identity that topologically generates G if and only if D contains an infinite suitable set for G .

Hofmann and Morris discovered the following fundamental result:

Theorem 1.9. ([13]; see also [14]) *Every compact group has a suitable set.*³

Remark 1.10. (i) *Theorem 1.6 implies the particular case of Theorem 1.9 for abelian groups.* Indeed, let G be a compact abelian group. If G is finite, then G is discrete, and so G itself is a suitable set for G . Assume now that G is infinite. By Theorem 1.6, G contains a qc-dense super-sequence S converging to 0. By Fact 1.7, S topologically generates G . According to Remark 1.8, $S \setminus \{0\}$ is a suitable set for G .

(ii) *A suitable set for a compact abelian group G need not be qc-dense in G .* Indeed, it is well-known that the group $\mathbb{T}^{\mathfrak{c}}$ is monothetic, that is, topologically generated by a singleton S . Clearly, S is a suitable set for $\mathbb{T}^{\mathfrak{c}}$. Since $w(S) \leq \omega < \mathfrak{c} = w(\mathbb{T}^{\mathfrak{c}})$, S cannot be qc-dense in $\mathbb{T}^{\mathfrak{c}}$ by Theorem 1.5.

(iii) It follows from item (ii) that *the particular case of Theorem 1.9 for abelian groups does not imply Theorem 1.6.*

2. RESULTS

Our first result is a particular version of Theorem 1.6 that characterizes connected compact abelian groups.

Theorem 2.1. *For an infinite compact abelian group G the following conditions are equivalent:*

- (i) *the arc component G_a of G contains a super-sequence converging to 0 that is qc-dense in G ;*
- (ii) *G is connected.*

In view of Fact 1.4, the implication (ii)→(i) of Theorem 2.1 yields Theorem 1.3 when G is compact. The general case of Theorem 1.3 easily follows from the compact case, see the proof in the end of Section 4. Therefore, as a by-product, our proof of Theorem 2.1 also provides an alternative short and self-contained proof of the theorem of Außenhofer.

Remark 2.2. In view of Theorems 1.6 and 2.1, as well as Außenhofer's result cited in footnote 2, the reader may wonder if every dense subgroup of a compact abelian group G contains a super-sequence converging to 0 that is qc-dense in G . The answer to this question is negative: *Every non-metrizable compact abelian group G contains a dense subgroup H such that no super-sequence $S \subseteq H$ is qc-dense in G .* Indeed, apply Theorem 1.2 to get a

³A “purely topological” proof of this result based on Michael's selection theorem can be found in [16].

dense subgroup H of G that does not determine G . If $S \subseteq H$ is a super-sequence, then (being compact) S cannot be qc-dense in G by Fact 1.4.

Our second result is a particular version of Theorem 1.9 that characterizes connected compact groups:

Theorem 2.3. *For an infinite compact group G the following conditions are equivalent:*

- (i) *the arc component G_a of G contains a suitable set for G ;*
- (ii) *G_a contains an infinite suitable set for G that is qc-dense in G ;*
- (iii) *G is connected.*

Remark 2.4. One cannot add the following item to the list of equivalent conditions in Theorem 2.3:

- (iv) *G_a contains a super-sequence converging to the identity that is qc-dense in G .*

Indeed, for every finite simple non-commutative group L the compact group $G = \mathbb{T} \times L$ has a sequence $S \subseteq G_a = \mathbb{T} \times \{e\}$ converging to the identity $(0, e)$ of G that is qc-dense in G , see Example 5.3. Since G is not connected, the implication (iv) $\rightarrow$ (iii) fails.

Remark 2.5. Let $\mathfrak{c}$ denote the cardinality of the continuum. *There exists a dense (connected, locally connected, countably compact) subgroup H of (the compact, connected abelian group) $G = \mathbb{T}^{2^{\mathfrak{c}}}$ such that H contains no suitable set for G .* Indeed, one can take as H the dense subgroup of G without a suitable set for H constructed in [11, Corollary 2.9]. Assume that $S \subseteq H$ is a suitable set for G . Then S is discrete and $S \cup \{0\}$ is closed in G . Since $S \cup \{0\} \subseteq H$, it follows that S is closed in H . Since S topologically generates G , it topologically generates H as well. Hence, S is a suitable set for H , a contradiction.

3. A QC-DENSE SUPER-SEQUENCE IN THE ARC COMPONENT OF $\hat{\mathbb{Q}}$

Our main result in this section is Lemma 3.4. It follows from the density of $\hat{\mathbb{Q}}_a$ in $\hat{\mathbb{Q}}$ and the general result of Außenhofer quoted in the footnote 2. However, Außenhofer’s proof relies on Arzela-Ascoli theorem and an inductive construction, so the qc-dense sequence she constructs in her proof is “generic”. To keep this manuscript self-contained, we provide a “constructive” example of a “concrete” qc-dense sequence in $\mathbb{Q}_a$.

The proof of the following fact is straightforward from the definition.

Fact 3.1. *Let G and H be topological groups and $\pi : H \rightarrow G$ a continuous surjective group homomorphism. If a subset E of H is qc-dense in H , then $\pi(E)$ is qc-dense in G .*

In the sequel $\mathbb{N}$ denotes the set of natural numbers.

Example 3.2. Let $T = \left\{ \frac{1}{2^n} : n \in \mathbb{N}, n \geq 1 \right\} \cup \{0\}$. The set $\varphi(T)$ is a qc-dense sequence in $\mathbb{T}$ converging to 0. Indeed, let $\chi \in \hat{\mathbb{T}}$ be a non-zero character. Then there exists $m \in \mathbb{Z} \setminus \{0\}$ such that $\chi(x) = mx$ for all $x \in \mathbb{T}$. Let $n = |m|$. Then $\frac{1}{2^n} \in T$ and so $x = \varphi\left(\frac{1}{2^n}\right) \in \varphi(T)$. Since $\chi(x) = mx = \varphi\left(\frac{m}{2^n}\right) = \varphi\left(\frac{1}{2}\right) \notin \mathbb{T}_+$, we have $\chi(\varphi(T)) \setminus \mathbb{T}_+ \neq \emptyset$. This proves that $\varphi(T)$ is qc-dense in $\mathbb{T}$.

For $g \in G$ the symbol $\langle g \rangle$ denotes the cyclic subgroup of G generated by g .

Lemma 3.3. *Let $\mathbb{P} = \{p_n : n \in \mathbb{N}\}$ be a faithful enumeration of the set $\mathbb{P}$ of prime numbers. Define*

$$H = \prod_{n \in \mathbb{N}} \mathbb{Z}_{p_n},$$

and let $v = \{1_{p_n}\}_{n \in \mathbb{N}} \in H$, where each 1_{p_n} is the identity of $\mathbb{Z}_{p_n}$. For $n \in \mathbb{N}$ define $k_n = (p_0 p_1 \dots p_{n-1})^n$. Then the set

$$(1) \quad S = \{mk_n v : n \in \mathbb{N}, m \leq k_{n+1}\} \cup \{0\} \subseteq \langle v \rangle$$

is a sequence converging to 0 that is qc-dense in H .

Proof. For $n \in \mathbb{N}$ define

$$(2) \quad W_n = k_n H = p_0^n \mathbb{Z}_{p_0} \times p_1^n \mathbb{Z}_{p_1} \times \dots \times p_{n-1}^n \mathbb{Z}_{p_{n-1}} \times \prod_{i=n}^{\infty} \mathbb{Z}_{p_i}.$$

(Note that $k_0 = 1$.) Then $\{W_n : n \in \mathbb{N}\}$ forms a base of H at 0 consisting of clopen subgroups. It is easy to see that each W_n may miss only finitely many members of S , so S is a sequence converging to 0 in H .

Let us show that S is qc-dense in H . Let $\chi \in \widehat{H}$ and $\chi \neq 0$. We need to prove that $\chi(S) \setminus \mathbb{T}_+ \neq \emptyset$. Being a continuous homomorphic image of the compact totally disconnected group H , $\chi(H)$ is a closed totally disconnected subgroup of $\mathbb{T}$. Therefore, $\chi(H)$ must be finite. Hence $\ker \chi$ is an open subgroup of H , and consequently it contains a subgroup W_n for some $n \in \mathbb{N}$. Without loss of generality we will assume that

$$(3) \quad n = \min\{m \in \mathbb{N} : W_m \subseteq \ker \chi\}.$$

Since $\ker \chi \neq H = W_0$ by our assumption, we have $n \geq 1$, and so $n-1 \in \mathbb{N}$.

Claim: $\chi(k_{n-1}v) \neq 0$.

Proof. Assume the contrary. Then $\chi \upharpoonright_{\langle k_{n-1}v \rangle} = 0$. Since $\langle v \rangle$ is dense in H and W_{n-1} is an open subset of H , it follows that $\langle v \rangle \cap W_{n-1} = \langle k_{n-1}v \rangle$ is dense in W_{n-1} . Now from $\chi \upharpoonright_{\langle k_{n-1}v \rangle} = 0$ and continuity of χ we conclude that $\chi \upharpoonright_{W_{n-1}} = 0$. This gives $W_{n-1} \subseteq \ker \chi$, in contradiction with (3). $\square$

Since $k_n v \in W_n \subseteq \ker \chi$ by (2) and (3), we have $k_n \chi(v) = \chi(k_n v) = 0$. That is, $\langle \chi(v) \rangle$ is a cyclic group of order at most k_n . Since $\chi(k_{n-1}v) = k_{n-1} \chi(v) \in \langle \chi(v) \rangle$, the order of the element $\chi(k_{n-1}v)$ of $\mathbb{T}$ is also at most k_n . Since $\chi(k_{n-1}v) \neq 0$ by claim, we can choose an integer $m \leq k_n$ such that $\chi(mk_{n-1}v) = m\chi(k_{n-1}v) \notin \mathbb{T}_+$. From (1) we conclude that $mk_{n-1}v \in S$, and so $\chi(S) \setminus \mathbb{T}_+ \neq \emptyset$. $\square$

An explicit qc-dense sequence in $\widehat{\mathbb{Q}}$ converging to 0 can be found in [10, Lemma 4.7]. However, that sequence is not contained in $\widehat{\mathbb{Q}}_a$. In our next lemma we produce a qc-dense sequence converging to 0 inside $\widehat{\mathbb{Q}}_a$.

Lemma 3.4. *$\widehat{\mathbb{Q}}_a$ contains a sequence converging to 0 that is qc-dense in $\widehat{\mathbb{Q}}$.*

Proof. We continue using notations from Lemma 3.3. Let $K = \mathbb{R} \times H$ and $u = (1, v) \in K$. Then the cyclic subgroup $\langle u \rangle$ of K is discrete and the quotient group $C = K/\langle u \rangle$ is isomorphic to $\widehat{\mathbb{Q}}$ [7, §2.1]. Therefore, it suffices to prove that C_a contains a sequence converging to 0 that is qc-dense in C .

Let $\pi : K \rightarrow C = K/\langle u \rangle$ be the quotient homomorphism. Since π is a local homeomorphism, every continuous map $f : [0, 1] \rightarrow C$ with $f(0) = 0_C$ can be lifted to a continuous map $\tilde{f} : [0, 1] \rightarrow K$ with $\tilde{f}(0) = 0_K$ and $\pi \circ \tilde{f} = f$. (A more general statement can be found in [15, Lemma 1].) Therefore, $C_a = \pi(K_a)$. Since H is zero-dimensional and $\mathbb{R} \times \{0\}$ is arcwise connected, one has $K_a = \mathbb{R} \times \{0\}$, and so $C_a = \pi(\mathbb{R} \times \{0\})$.

Define $N = \pi(\{0\} \times H)$, and let $f : C \rightarrow C/N$ be the quotient homomorphism. By Lemma 3.3 and Fact 3.1, there exists a converging to 0 sequence S' in the subgroup $\langle \pi(0, v) \rangle$ of N such that S' is qc-dense in N . As

$$\pi(0, v) = \pi((-1, 0) + (1, v)) = \pi(-1, 0) + \pi(u) = -\pi(1, 0) \in \pi(\mathbb{R} \times \{0\}) = C_a,$$

one has $S' \subseteq \pi(\langle (0, v) \rangle) \subseteq C_a$.

With $T = \left\{ \frac{1}{2^n} : n \in \mathbb{N}, n \geq 1 \right\}$ define $S'' = \pi(T \times \{0\}) \subseteq \pi(\mathbb{R} \times \{0\}) = C_a$. Clearly, S'' is a sequence converging to 0. Since $C/N \cong K/(\mathbb{Z} \times H) \cong \mathbb{T}$ and the composed isomorphism $C/N \rightarrow \mathbb{T}$ sends $f(S'')$ to $\varphi(T)$, from Example 3.2 we conclude that $f(S'')$ is qc-dense in C/N .

Since S' and S'' are sequences converging to 0 in C , so is $X = S' \cup S''$. By our construction, $X \subseteq C_a$. So it remains only to prove that X is qc-dense in C . Suppose that $\chi \in \widehat{C}$ and $\chi(X) \subseteq \mathbb{T}_+$. Since $\chi \upharpoonright_N \in \widehat{N}$, $\chi \upharpoonright_N (S') = \chi(S') \subseteq \widehat{\chi(X)} \subseteq \mathbb{T}_+$ and S' is qc-dense in N , we have $\chi \upharpoonright_N = 0$. Therefore, $\chi = \xi \circ f$ for some $\xi \in \widehat{C/N}$. In particular, $\xi(f(S'')) \subseteq \xi(f(X)) = \chi(X) \subseteq \mathbb{T}_+$. Since $f(S'')$ is qc-dense in C/N , it follows that $\xi = 0$. This gives $\chi = 0$. Therefore, X is qc-dense in C . $\square$

4. PROOF OF THEOREMS 2.1 AND 1.3

The following definition is an adaptation to the abelian case of [9, Definition 4.5]:

Definition 4.1. Let $\{G_i : i \in I\}$ be a family of abelian topological groups. For every $i \in I$ let X_i be a subset of G_i . Identifying each G_i with a subgroup of the direct product $G = \prod_{i \in I} G_i$ in the obvious way, define $X = \bigcup_{i \in I} X_i \cup \{0\}$, where 0 is the zero element of H . We will call X the *fan* of the family $\{X_i : i \in I\}$ and will denote it by $\text{fan}_{i \in I}(X_i, G_i)$.

The proof of the following lemma is straightforward.

Lemma 4.2. [10, Lemmas 4.3 and 4.4] *Let $\{G_i : i \in I\}$ be a family of abelian topological groups, and let $G = \prod_{i \in I} G_i$. For every $i \in I$ let X_i be a subset of G_i , and let $X = \text{fan}_{i \in I}(X_i, G_i)$. Then:*

- (i) *if X_i is a sequence converging to 0 in G_i , then X is a super-sequence in G converging to 0.*
- (ii) *if X_i is a qc-dense subset of G_i for each $i \in I$, then X is qc-dense in G .*

Lemma 4.3. *For every cardinal κ there exists a super-sequence $S \subseteq (\widehat{\mathbb{Q}}^\kappa)_a$ converging to 0 that is qc-dense in $\widehat{\mathbb{Q}}^\kappa$.*

Proof. Write $\widehat{\mathbb{Q}}^\kappa$ as $\widehat{\mathbb{Q}}^\kappa = \prod_{\alpha < \kappa} G_\alpha$, where G_α is the α 's copy of $\widehat{\mathbb{Q}}$. By Lemma 3.4, for every $\alpha \in \kappa$ there is a sequence S_α in $(G_\alpha)_a$ converging to 0 that is qc-dense in G_α . By Lemma 4.2(i), $S = \text{fan}_{\alpha \in \kappa}(X_\alpha, G_\alpha)$ is a super-sequence in $\widehat{\mathbb{Q}}^\kappa$ converging to 0. By Lemma 4.2(ii), S is qc-dense in $\widehat{\mathbb{Q}}^\kappa$. Finally, note that $S \subseteq \bigoplus_{\alpha \in \kappa} (G_\alpha)_a \subseteq (\widehat{\mathbb{Q}}^\kappa)_a$. $\square$

Proof of Theorem 2.1. (i)→(ii) Let $S \subseteq G_a$ be a super-sequence that is qc-dense in G . Then the subgroup H of G generated by S is dense in G by Fact 1.7. Since $H \subseteq G_a$, it follows that G_a is dense in G as well. Thus G is connected.

(ii)→(i) There exists a continuous surjective homomorphism $\pi : \hat{\mathbb{Q}}^\kappa \rightarrow G$ for some cardinal κ (see, for example, the proof of [10, Theorem 3.3]). Let S be as in the conclusion of Lemma 4.3. Since S is qc-dense in $\hat{\mathbb{Q}}^\kappa$, $\pi(S)$ is qc-dense in G by Fact 3.1. Since a finite set cannot be qc-dense in an infinite compact group ([1]; this also follows from Theorem 1.5), $\pi(S)$ must be infinite. Since S is a super-sequence converging to 0 such that $\pi(S)$ is infinite, $\pi(S)$ must be a super-sequence converging to 0 by [9, Fact 4.3] (see [16, Fact 12] for the proof). Finally note that $\pi(S) \subseteq \pi(\hat{\mathbb{Q}}_a^\kappa) \subseteq G_a$. $\square$

Proof of Theorem 1.3. We have $G = \mathbb{R}^n \times K$, where K is a compact connected group [8]. Since $\mathbb{R}^n$ is arcwise connected, one has $G_a = \mathbb{R}^n \times K_a$. From Theorem 2.1 and Fact 1.4 we conclude that K_a determines K . Hence $G_a = \mathbb{R}^n \times K_a$ determines $G = \mathbb{R}^n \times K$. $\square$

5. PROOF OF THEOREM 2.3

In the sequel we denote by H' the commutator subgroup of a group H .

Our next lemma shows that qc-density can be essentially studied in the abelian context.

Lemma 5.1. *Let H be a topological group, and let G denote the quotient $H/\overline{H'}$, where $\overline{H'}$ is the closure of H' in H . Let $\pi : H \rightarrow G$ denote the canonical map. Then a subset E of H is qc-dense in H if and only if $\pi(E)$ is qc-dense in G .*

Proof. The “only if” part follows from Lemma 3.1. To prove the “if” part, assume that $\pi(E)$ is qc-dense in G , and let $\chi : H \rightarrow \mathbb{T}$ be a continuous homomorphism such that $\chi(E) \subseteq \mathbb{T}_+$. Since $\mathbb{T}$ is abelian and Hausdorff, $\overline{H'} \subseteq \ker \chi$, so $\chi = \xi \circ \pi$ for some character $\xi : G \rightarrow \mathbb{T}$. Since $\xi(\pi(E)) = \chi(E) \subseteq \mathbb{T}_+$ and $\pi(E)$ is qc-dense in G , we conclude that ξ is trivial, and so χ is trivial too. This proves that E is qc-dense in H . $\square$

Recall that a group L is called *perfect* if $L' = L$.

Corollary 5.2. *Let L be a perfect topological group, G an abelian topological group and $H = G \times L$. Then a subset E of G is qc-dense in G if and only if the subset $E \times \{e_L\}$ of the group H is qc-dense in H .*

Proof. Since $H' = \{0_G\} \times L$, we have $H' = \overline{H'}$ and $G \cong H/H' = H/\overline{H'}$. Now the conclusion of our corollary follows from Lemma 5.1 applied to the projection $\pi : H \rightarrow G$. $\square$

Example 5.3. Let $T = \left\{ \frac{1}{2^n} : n \in \mathbb{N}, n \geq 1 \right\} \cup \{0\}$. Assume that L is a finite simple non-abelian group and $G = \mathbb{T} \times L$. Then $S = \varphi(T) \times \{e_L\}$ is a qc-dense sequence in G converging to e_G . Indeed, since all simple non-abelian groups are perfect, this follows from Example 3.2 and Corollary 5.2.

Proof of Theorem 2.3. (i)→(iii) Let $S \subseteq G_a$ be a suitable set for G . Then the subgroup H of G generated by S is dense in G . Since $H \subseteq G_a$, it follows that G_a is dense in G as well. Thus G is connected.

(iii)→(ii) According to [9, Theorem 3.3], there exist a cardinal κ and a continuous surjective group homomorphism $\pi : \hat{\mathbb{Q}}^\kappa \times L \rightarrow G$, where L is a direct product of simple Lie groups. In particular, L is perfect. By Lemma 4.3, there exists a super-sequence E in $(\hat{\mathbb{Q}}^\kappa)_a$ converging to 0 that is qc-dense in $\hat{\mathbb{Q}}^\kappa$. By Corollary 5.2, $S' = E \times \{e_L\}$ is a converging to $(0, e_L)$

super-sequence in $(\widehat{\mathbb{Q}}^\kappa)_a \times \{e_L\}$ that is qc-dense in $\widehat{\mathbb{Q}}^\kappa \times L$. Since S' is qc-dense in $\widehat{\mathbb{Q}}^\kappa \times \{e\}$, it topologically generates $\widehat{\mathbb{Q}}^\kappa \times \{e\}$ by Fact 1.7. By Theorem 1.9 and Remark 1.8, there exists a super-sequence $S'' \subseteq \{0\} \times L$ converging to $(0, e_L)$ that topologically generates $\{0\} \times L$. Then $S = S' \cup S''$ is a super-sequence converging to $(0, e_L)$ that is qc-dense in $\widehat{\mathbb{Q}}^\kappa \times L$ and topologically generates $\widehat{\mathbb{Q}}^\kappa \times L$. Therefore, $X = \pi(S)$ is a super-sequence (by [9, Fact 4.3]; see [16, Fact 12] for the proof) that is qc-dense in G (by Fact 3.1) and topologically generates G . Since L is arcwise connected and $S \subseteq (\widehat{\mathbb{Q}}^\kappa)_a \times L$, we have $X = \pi(S) \subseteq \pi((\widehat{\mathbb{Q}}^\kappa)_a \times L) \subseteq G_a$. Applying Remark 1.8, we conclude that $X \setminus \{e\}$ is a suitable set for G . Since X is qc-dense in G , so is $X \setminus \{e\}$. If X is infinite, we are done.

Assume now that X is finite. Since S' is qc-dense in $\widehat{\mathbb{Q}}^\kappa \times \{e\}$, $\pi(S')$ is qc-dense in $K = \pi(\widehat{\mathbb{Q}}^\kappa \times \{e\})$ by Fact 3.1. Since $\pi(S') \subseteq \pi(S) = X$, the set $\pi(S')$ is finite. From Theorem 1.5 we conclude that the (compact abelian) group K must be finite as well. Being a continuous image of the connected group $\widehat{\mathbb{Q}}^\kappa \times \{e\}$, the group K is connected. Hence, K is trivial, and so $G = \pi(\widehat{\mathbb{Q}}^\kappa \times L) = \pi(\{0\} \times L) \subseteq G_a \subseteq G$ because L is pathwise connected. This yields $G = G_a$.

Let $f : [0, 1] \rightarrow G_a$ be a continuous map such that $f(0) = e_G$ and $f(1) \neq e_G$. Define $c = \inf\{t \in [0, 1] : f(t) \neq e_G\}$. Then $f(c) = e_G$ by continuity of f and the choice of c . For every $n \in \mathbb{N}$ choose $c_n \in [0, 1]$ such that $c < c_n < c + 1/n$ and $f(c_n) \neq e_G$. Since $\{c_n : n \in \mathbb{N}\}$ converges to c and f is continuous, the sequence $X_0 = \{f(c_n) : n \in \mathbb{N}\}$ converges to $f(c) = e_G$. Since $f(c_n) \neq e_G$ for every $n \in \mathbb{N}$, we conclude that X_0 is infinite. Now $X' = X \cup X_0 \cup \{e_G\}$ is an infinite super-sequence converging to e_G , and so $X' \setminus \{e_G\} \subseteq G_a$ is an infinite suitable set for G by Remark 1.8. Since X is qc-dense in G and topologically generates G , X' has the same properties.

The implication (ii) $\rightarrow$ (i) is trivial. $\square$

REFERENCES

- [1] L. Außenhofer, *Contributions to the duality theory of abelian topological groups and to the theory of nuclear groups*. Diss. Math. CCCLXXXIV. Warsaw, 1999.
- [2] L. Außenhofer, A duality property of an uncountable product of $\mathbb{Z}$, *Math. Z.* 257 (2007) 231–237.
- [3] L. Außenhofer, On the arc component of a locally compact abelian group, *Math. Z.* 257 (2007) 239–250.
- [4] M. J. Chasco, Pontryagin duality for metrizable groups, *Arch. Math. (Basel)* 70 (1998) 22–28.
- [5] W. W. Comfort, S. U. Raczkowski and F. J. Trigos-Arrieta, The dual group of a dense subgroup, *Czechoslovak Math. J.* 54 (129) (2004) 509–533.
- [6] D. Dikranjan and L. De Leo, Countably infinite quasi-convex sets in some locally compact abelian groups, submitted.
- [7] D. Dikranjan and C. Milan, Dualities of locally compact modules over the rationals, *J. Algebra* 256 (2002) 433–466.
- [8] D. Dikranjan, Iv. Prodanov and L. Stoyanov, *Topological Groups: Characters, Dualities and Minimal Group Topologies*, Pure and Applied Mathematics, vol. 130, Marcel Dekker Inc., New York-Basel (1989).
- [9] D. Dikranjan and D. Shakhmatov, Weight of closed subsets topologically generating a compact group, *Math. Nachr.* 280 (2007) 505–522.
- [10] D. Dikranjan and D. Shakhmatov, Quasi-convex density and determining subgroups of compact abelian groups, submitted (available as ArXiv preprint no. arXiv:0807.3846v3).
- [11] D. Dikranjan, M. Tkačenko, and V. Tkachuk, Some topological groups with and some without suitable sets, *Topol. Appl.* 98 (1999) 131–148.
- [12] S. Hernández, S. Macario and F. J. Trigos-Arrieta, Uncountable products of determined groups need not be determined, *J. Math. Anal. Appl.* 348 (2008) 834–842.
- [13] K.-H. Hofmann and S. A. Morris, Weight and c , *J. Pure Appl. Algebra* 68 (1990) 181–194.

- [14] K.-H. Hofmann and S. A. Morris, *The structure of compact groups. A primer for the student—a handbook for the expert*, de Gruyter Studies in Mathematics, 25, Walter de Gruyter & Co., Berlin, 1998.
- [15] N. W. Rickert, *Arcs in locally compact groups*, *Math. Ann.* 172 (1967) 222–228.
- [16] D. Shakhmatov, *Building suitable sets for locally compact groups by means of continuous selections*, *Topol. Appl.* (2008), doi:10.1016/j.topol.2008.12.009 (available as ArXiv preprint no. arXiv:0812.0489v2).

(Dikran Dikranjan) UNIVERSITÀ DI UDINE, DIPARTIMENTO DI MATEMATICA E INFORMATICA, VIA DELLE SCIENZE, 206 - 33100 UDINE, ITALY

E-mail address: `dikran.dikranjan@dimi.uniud.it`

(Dmitri Shakhmatov) GRADUATE SCHOOL OF SCIENCE AND ENGINEERING, DIVISION OF MATHEMATICS, PHYSICS AND EARTH SCIENCES, EHIME UNIVERSITY, MATSUYAMA 790-8577, JAPAN

E-mail address: `dmitri@dpc.ehime-u.ac.jp`